



Introduction to neutron-induced reactions and the R-matrix formalism

Frank Gunsing

CEA/Saclay

DSM / Irfu / SPhN

F - 91911 Gif-sur-Yvette, France

gunsing@cea.fr

Currently on leave at

CERN

CH-1211 Geneva 23

Switzerland





CV

Education

- 1/6/2005 **Habilitation à Diriger des Recherches (HDR)**, *Université Paris VII (France)*.
- 1991 - 1995 **PhD in Physics**, *Delft University of Technology (The Netherlands)*.
- 1988 - 1989 **Maîtrise de Physique et ses Applications**, *Université Paris VII (France)*.
- 1985 - 1991 **Engineer in Physical Sciences**, *Delft University of Technology (The Netherlands)*.



Work Experience

- 1996 Jan - present **Staff physicist**, *CEA Saclay, France. DSM-Irfu, Nuclear Physics Division.*
 - Research activities: experimental nuclear physics, neutron-nucleus reactions, surrogate reactions, compound nucleus spectroscopy, nuclear data, detector development
Experimental activities currently at: JRC-IRMM-Geel, OCL-Oslo, n_TOF-CERN
 - Implication in European Framework Projects FP5, FP6, FP7
 - Scientific and budget responsible
 - Supervision postdocs, PhD and master students
- 1995 July-Dec **Postdoc**, *CEA Saclay, France*, Study of neutron induced reactions for nuclear waste transmutation.
- 1995 Jan-June **Postdoc**, *Joint Research Centre - IRMM, Geel, Belgium*, Study of neutron induced reactions for nuclear waste transmutation.
- 1991 - 1994 **PhD student**, *Joint Research Centre - IRMM, Geel, Belgium*, Phd work on neutron resonance spin measurements for parity violation.
- 1989 - 1991 **master student**, *Low Temperature Laboratory, Delft University of Technology, The Netherlands*, Development of RuO₂ bolometers.

Scientific and Editorial Committees



Neutron induced reactions





Neutron-nucleus reactions

Reaction:

- $X + a \rightarrow Y + b$
- $X(a,b)Y$
- $X(a,b)$

Examples of equivalent notations:



Reaction cross section σ , expressed in barns, $1 \text{ b} = 10^{-28} \text{ m}^2$


Neutron induced nuclear reactions:

- elastic scattering (n,n)
- inelastic scattering (n,n')
- capture (n,γ)
- fission (n,f)
- particle emission (n,α) , (n,p) , (n,xn)

Total cross section σ_{tot} : sum of all reactions

Neutron-nucleus reactions

Reaction: $X + a \rightarrow Y + b$
 $X(a,b)Y$



Cross section:

function of the kinetic energy of the particle a $\sigma(E_a) = \int \int \frac{d^2\sigma(E_a, E_b, \Omega)}{dE_b d\Omega} dE_b d\Omega$


Differential cross section:

function of the kinetic energy of the particle a and function of the kinetic energy **or** the angle of the particle b $\frac{d\sigma(E_a, E_b)}{dE_b}$ $\frac{d\sigma(E_a, \Omega)}{d\Omega}$


Double differential cross section:

function of the kinetic energy of the particle a and function of the kinetic energy **and** the angle of the particle b $\frac{d^2\sigma(E_a, E_b, \Omega)}{dE_b d\Omega}$


Cross sections σ_T , σ_γ , σ_n et σ_f




Neutron fluxes and cross sections




Neutron fluxes and cross sections




Neutron fluxes and cross sections





Neutron fluxes and cross sections





Neutron induced reaction cross sections





Neutron fluxes and cross sections



Neutron fluxes and cross sections



Interference of $\sigma_{\text{potential}}$ and σ_n



transmission
 $T = \exp(-n \cdot \sigma_T)$
 $0 < T < 1$

Classical – Quantum Physics

Classical physics

- particles, Newton's law of motion
- electromagnetic waves, Maxwell's laws of electromagnetism

Quantum physics

- particles (momentum) and waves (wavelength) are different descriptions of the same thing. Related by Planck's constant h .

$$\text{De Broglie wavelength: } \lambda = \frac{h}{p}$$

From 1900, observations of electrons, photons behaving as particles or waves in different experiments (black body radiation, photo-electric effect, crystal diffraction).

Probability of a particle being at time t , having position x is related to a “wave function”.

Probability (Born interpretation): $\psi^* \psi$




The wave function is a solution of the Schrödinger equation (postulate).

The hydrogen atom


Hydrogen atom

- quantum system of one proton, one electron
- the system can be in well-defined energy states (electron orbits).
- transitions between these states can be observed as electromagnetic radiation
- Observed: energy states: $E_n = -13.6/n^2$ eV, with $n=1$ the ground state
- wavelengths observed corresponding to transitions between these states ($\Delta E=hc/\lambda$)


The hydrogen atom – Bohr model





The hydrogen atom – Bohr model



The hydrogen atom – Bohr model






Discrete states of the hydrogen atom



Discrete states of the hydrogen atom

Hydrogen atom

- quantum system of one proton, one electron
- the system can be in well-defined energy states (electron orbits).
- transitions between these states can be observed as electromagnetic radiation
- energy states: $E_n = -13.6/n^2$ eV, with $n=1$ the ground state
- wavelengths observed corresponding to transitions between these states ($\Delta E=hc/\lambda$)




The Schrödinger equation

Time-independent, for a spinless, onedimensional particle:

wave function

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

potential energy



Solutions: ψ, E

$\psi^* \psi$ Interpreted as probability

The Schrödinger equation

Time-independent. Equation for a spinless, onedimensional particle:

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V(x) \right] \psi(x) = E\psi(x)$$

$$\hat{H}\psi(x) = E\psi(x)$$



Hamiltonian

Quantum system: the infinite well

Solve Schrödinger equation,
for a spinless, onedimensional particle

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E(x)$$


Probability $\psi^*\psi$

Example: The infinite well
for a spinless, onedimensional particle:

$$V(x) = 0 \quad \text{for } 0 < x < a$$

$$V(x) = \infty \quad \text{elsewhere}$$

Solution:



Quantum system: the infinite well

Solve Schrödinger equation,
for a spinless, onedimensional particle

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E(x)$$

Probability $\psi^* \psi$

Example: The infinite well

for a spinless, onedimensional particle:


$$V(x) = 0 \quad \text{for } 0 < x < a$$

$$V(x) = \infty \quad \text{elsewhere}$$

Solution:

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin \frac{n\pi x}{a}$$

$$E_n = \frac{\hbar^2 \pi^2}{2ma^2} n^2$$



Quantum system: the infinite well

Solve Schrödinger equation,
for a spinless, onedimensional particle

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E(x)$$

Probability $\psi^* \psi$

Example: The infinite well

for a spinless, onedimensional particle:


$$V(x) = 0 \quad \text{for } 0 < x < a$$

$$V(x) = \infty \quad \text{elsewhere}$$

Solution:

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin \frac{n\pi x}{a}$$

$$E_n = \frac{\hbar^2 \pi^2}{2ma^2} n^2$$



Quantum system: the finite well

Solve Schrödinger equation in two regions:


- inside and outside the well
- normalize solutions to match value and derivative at borders $x=0$ and $x=a$

Now the wave function exists also outside the well at $x < 0$ and $x > a$

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

$$0 < x < a \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V_0\psi(x) = E\psi(x)$$

$$x < 0, x > a \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} = E\psi(x)$$



Quantum system: the finite well

Solve Schrödinger equation in two regions:


- inside and outside the well
- normalize solutions to match value and derivative at borders $x=0$ and $x=a$

Now the wave function exists also outside the well at $x < 0$ and $x > a$

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

$$0 < x < a \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V_0\psi(x) = E\psi(x)$$

$$x < 0, x > a \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} = E\psi(x)$$



Match value and derivative

Quantum system: the finite well


Solve Schrödinger equation in two regions:

- inside and outside the well
- normalize solutions to match value and derivative at borders $x=0$ and $x=a$

Now the wave function exists also outside the well at $x < 0$ and $x > a$

In general, a generic state can be written as a linear expansion of its eigenstates:


$$\psi(x) = \sum_k c_k \psi_k(x)$$



Quantum system: the potential barrier

Solve Schrödinger equation in three regions:


- free travelling particle of energy E
- inside and outside the well
- normalize solutions to match value and derivative and borders $x=0$ and $x=a$
- **transmission and reflection**



Quantum system: the potential barrier

Solve Schrödinger equation in three regions:

- free travelling particle of energy E
- inside and outside the well
- normalize solutions to match value and derivative and borders $x=0$ and $x=a$
- **transmission and reflection**



Quantum system: the potential barrier

Solve Schrödinger equation in three regions:


- free travelling particle of energy E
- inside and outside the well
- normalize solutions to match value and derivative and borders $x=0$ and $x=a$
- **transmission and reflection**

$$\psi_1(x) = Ae^{ik(x)x} + Be^{-ik(x)x}$$

$$\psi_2(x) = Ce^{ik(x)x} + De^{-ik(x)x}$$

$$\psi_3(x) = Ee^{ik(x)x} + Fe^{-ik(x)x}$$

$$k(x) = \sqrt{2m(E - V_0)/\hbar^2}$$



Quantum system: the potential barrier

Solve Schrödinger equation in three regions:


- free travelling particle of energy E
- inside and outside the well
- normalize solutions to match value and derivative and borders $x=0$ and $x=a$
- **transmission and reflection**

$$\psi_1(x) = Ae^{ik(x)x} + Be^{-ik(x)x}$$

$$\psi_2(x) = Ce^{ik(x)x} + De^{-ik(x)x}$$

$$\psi_3(x) = Ee^{ik(x)x} + Fe^{-ik(x)x}$$

$$k(x) = \sqrt{2m(E - V_0)/\hbar^2}$$



Quantum system: the potential barrier

Solve Schrödinger equation in three regions:

- free travelling particle of energy E
- inside and outside the well
- normalize solutions to match value and derivative at borders $x=0$ and $x=a$
- **transmission and reflection**


$$\psi_1(x) = Ae^{ik(x)x} + Be^{-ik(x)x}$$

$$\psi_2(x) = Ce^{ik(x)x} + De^{-ik(x)x}$$

$$\psi_3(x) = Ee^{ik(x)x} + Fe^{-ik(x)x}$$

$$k(x) = \sqrt{2m(E - V_0)/\hbar^2}$$

$$j = \frac{\hbar}{2mi} (\psi^* \nabla \psi - \nabla \psi^* \psi)$$



Other useful exercises in 1D:

- barrier potential
- finite potential well
- harmonic oscillator

More complicated in 3D, $V=V(r)$, more particles, degeneracy:

- cartesian well
- spherical well
- harmonic oscillator
- realistic potentials (Wood-Saxon),

→ No analytical solution possible,
numerical solutions

Apply to real quantum systems:
atoms (hydrogen) but also to nuclei.

Other useful exercises in 1D:

- barrier potential
- finite potential well
- harmonic oscillator

More complicated in 3D, $V=V(r)$, degeneracy:

- cartesian well
- spherical well
- harmonic oscillator
- realistic potentials (Wood-Saxon),





→ No analytical solution possible,
numerical solutions

Apply to real quantum systems:
atoms (hydrogen) but also to nuclei.

A nucleus is a quantum system of
nucleons (protons and neutrons), bound
together by the strong force.




The nucleus as a quantum system, shell model



The nucleus as a quantum system

shell model representation:
configuration of nucleons in their potential



level scheme representation:
excited states of a nucleus
(shell model and other states)



The nucleus as a quantum system

shell model representation:
configuration of nucleons in their potential


level scheme representation:
excited states of a nucleus
(shell model and other states)



The nucleus as a quantum system

shell model representation:
configuration of nucleons in their potential


level scheme representation:
excited states of a nucleus
(shell model and other states)




The nucleus as a quantum system

shell model representation:
configuration of nucleons in their potential


level scheme representation:
excited states of a nucleus
(shell model and other states)




→ nuclear state




The nucleus as a quantum system




The nucleus as a quantum system




The nucleus as a quantum system





Nuclear levels




Compound neutron-nucleus reactions



Compound neutron-nucleus reactions



Neutron induced reactions: Chart of nuclides



Neutron induced reactions: Chart of nuclides




Chart of nuclides








Chart of nuclides



Neutron separation energy



Fission of $^{235}\text{U}+n$ et $^{238}\text{U}+n$



↑
excitation energy

Decay of a nuclear state

state with a life time τ :



$$\Psi(t) = \Psi_0 e^{-iE_0 t / \hbar} e^{-t / 2\tau}$$

definition (Heisenberg):


$$\Gamma = \frac{\hbar}{\tau}$$

Fourier transform gives energy profile:

$$I(E) = \frac{\Gamma / 2\pi}{(E - E_0)^2 + \Gamma^2 / 4}$$



Neutron-nucleus reactions



Conservation of probability density:

$$\sigma(\Omega) = \frac{r^2 j_{\text{out}}(r, \Omega)}{j_{\text{inc}}}$$

Solve Schrödinger equation of system to get cross sections.
Shape of wave functions of in- and outgoing particles are known,
potential is unknown. Two approaches:


- calculate potential (optical model calculations, smooth cross section)
- use eigenstates (R-matrix, resonances)

R-matrix formalism


partial incoming wave functions: \mathcal{I}_c
 partial outgoing wave functions: $\mathcal{O}_{c'}$
 related by collision matrix: $U_{cc'}$

cross section:

$$\sigma_{cc'} = \pi \lambda_c^2 |\delta_{c'c} - U_{c'c}|^2$$



Find the wave functions



$r > a_c$ external region

$r < a_c$ internal region

$r = a_c$ match value and derivate of Ψ

$$\left[\frac{d^2}{dr^2} - \frac{\ell(\ell + 1)}{r^2} - \frac{2m_c}{\hbar^2}(V - E) \right] rR(r) = 0$$

External region: **easy**, solve Schrödinger equation

central force, separate radial and angular parts.

$$\psi(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi)$$

solution: solve Schrödinger equation of relative motion:

- Coulomb functions
- special case of neutron particles (neutrons): fonctions de Bessel


Internal region: **very difficult**, Schrödinger equation cannot be solved directly

solution: expand the wave function as a linear combination of its eigenstates.


using the **R-matrix:**

$$R_{cc'} = \sum_{\lambda} \frac{\gamma_{\lambda c} \gamma_{\lambda c'}}{E_{\lambda} - E}$$


The R-matrix formalism



The R-matrix formalism



The R-matrix formalism



$$\Psi = \Psi(R_{cc'})$$

$$R_{cc'} = \sum_{\lambda} \frac{\gamma_{\lambda c} \gamma_{\lambda c'}}{E_{\lambda} - E}$$

$$\Psi = \sum y_c \mathcal{I}_c + \sum x_{c'} \mathcal{O}'_{c'}$$

$$x_{c'} \equiv^c - \sum U_{c'c'} y_c$$

$$\mathcal{I}_c = I_c r^{-\epsilon_1} \varphi_c i^l Y_{m_\ell}^l(\theta, \phi) / \sqrt{v_c}$$

$$\mathcal{O}_c = O_c r^{-1} \varphi_c i^l Y_{m_\ell}^l(\theta, \phi) / \sqrt{v_c}$$

The R-matrix formalism

The wave function of the system is a superposition of incoming and outgoing waves:

$$\Psi = \sum_c y_c \mathcal{I}_c + \sum_{c'} x_{c'} \mathcal{O}'_c$$

Incoming and outgoing wavefunctions have form:

$$\mathcal{I}_c = I_c r^{-1} \varphi_c i^l Y_{m_\ell}^\ell(\theta, \phi) / \sqrt{v_c}$$

$$\mathcal{O}_c = O_c r^{-1} \varphi_c i^l Y_{m_\ell}^\ell(\theta, \phi) / \sqrt{v_c}$$

The physical interaction is included in the collision matrix \mathbf{U} :

$$x_{c'} \equiv - \sum_c U_{c'c} y_c$$

The wave function depends on the R-matrix, which depends on the widths and levels of the eigenstates.

$$\Psi = \Psi(R_{cc'})$$

$$R_{cc'} = \sum_\lambda \frac{\gamma_{\lambda c} \gamma_{\lambda c'}}{E_\lambda - E}$$

The R-matrix formalism

The relation between the R-matrix and the collision matrix:

$$\mathbf{U} = \mathbf{\Omega} \mathbf{P}^{1/2} [\mathbf{1} - \mathbf{R}(\mathbf{L} - \mathbf{B})]^{-1} [\mathbf{1} - \mathbf{R}(\mathbf{L}^* - \mathbf{B})] \mathbf{P}^{-1/2} \mathbf{\Omega}$$

$$\text{with: } L_c = S_c + iP_c = \left(\frac{\rho}{O_c} \frac{dO_c}{d\rho} \right)_{r=a_c}$$

The relation between the collision matrix and cross sections:


$$\text{channel to one other channel: } \sigma_{cc'} = \pi \lambda_c^2 |\delta_{c'c} - U_{c'c}|^2$$

$$\text{channel to any other channel: } \sigma_{cr} = \pi \lambda_c^2 (1 - |U_{cc}|^2)$$


$$\text{channel to same channel: } \sigma_{cc} = \pi \lambda_c^2 |1 - U_{cc}|^2$$

$$\text{channel to any channel (total): } \sigma_{c,T} = \sigma_c = 2\pi \lambda_c^2 (1 - \text{Re} U_{cc})$$

The R-matrix formalism



The R-matrix formalism



The R-matrix formalism

The Breit-Wigner Single Level approximation:

total cross section:

$$\sigma_c = \pi \lambda_c^2 g_c \left(4 \sin^2 \phi_c + \frac{\Gamma_\lambda \Gamma_{\lambda c} \cos 2\phi_c + 2(E - E_\lambda - \Delta_\lambda) \Gamma_{\lambda c} \sin 2\phi_c}{(E - E_\lambda - \Delta_\lambda)^2 + \Gamma_\lambda^2/4} \right)$$

neutron channel: $c = n$

only capture, scattering, fission: $\Gamma_\lambda = \Gamma = \Gamma_n + \Gamma_\gamma + \Gamma_f$

other approximations: $\ell = 0$ $\cos \phi_c = 1$ $\sin \phi_c = \rho = ka_c$ $\Delta_\lambda = 0$

total cross section:

$$\sigma_T(E) = \overbrace{4\pi R'^2}^{\text{potential}} + \pi \lambda^2 g \left(\frac{\overbrace{4\Gamma_n(E - E_0)R'/\lambda}^{\text{interference}} + \overbrace{\Gamma_n^2}^{\text{elastic}} + \overbrace{\Gamma_n\Gamma_\gamma}^{\text{capture}} + \overbrace{\Gamma_n\Gamma_f}^{\text{fission}}}{\underbrace{(E - E_0)^2 + (\Gamma_n + \Gamma_\gamma + \Gamma_f)^2/4}_{\text{total width}}} \right)$$

The R-matrix formalism

The Reich-Moore approximation:

Use the fact that there are many photon channels, with Gaussian distributed amplitudes with zero mean:

$$\langle \gamma_{\lambda c} \gamma_{\mu c} \rangle = \gamma_{\lambda c}^2 \delta_{\lambda \mu}$$

The sum over the amplitudes of the photon channels becomes then:

$$\sum_{c \in \text{photon}} \gamma_{\lambda c} \gamma_{\mu c} = \sum_{c \in \text{photon}} \gamma_{\lambda c}^2 \delta_{\lambda \mu} = \Gamma_{\lambda \gamma} \delta_{\lambda \mu}$$

Then photon channels can be eliminated in the R-matrix:

$$R_{cc'} = \sum_{\lambda} \frac{\gamma_{\lambda c} \gamma_{\lambda c'}}{E_{\lambda} - E - i\Gamma_{\lambda \gamma}/2} \quad c \notin \text{photon}$$

Average cross sections

The relation between the energy averaged collision matrix and energy averaged cross sections:


average scattering:	$\overline{\sigma_{cc}} = \pi \lambda_c^2 g_c \overline{ 1 - U_{cc} ^2}$
shape elastic (potential)	$\overline{\sigma_{cc}^{se}} = \pi \lambda_c^2 g_c \overline{ 1 - \overline{U_{cc}} ^2}$
compound elastic	$\overline{\sigma_{cc}^{ce}} = \pi \lambda_c^2 g_c \left(\overline{ U_{cc} ^2} - \overline{U_{cc}} ^2 \right)$
average any reaction	$\overline{\sigma_{cr}} = \pi \lambda_c^2 g_c (1 - \overline{ U_{cc} ^2})$
average total	$\overline{\sigma_{c,T}} = 2\pi \lambda_c^2 g_c (1 - \text{Re} \overline{U_{cc}})$
average single reaction	$\overline{\sigma_{cc'}} = \pi \lambda_c^2 g_c \overline{ \delta_{cc'} - U_{cc'} ^2}$
average compound nucleus formation	$\overline{\sigma_c} = \pi \lambda_c^2 g_c (1 - \overline{ U_{cc} ^2})$

Average cross sections


- From optical model calculations one can calculate $\overline{U_{cc}}$ but not $|\overline{U_{cc}}|^2$
- Therefore, only $\overline{\sigma_{c,T}}$, $\overline{\sigma_{cc}^{se}}$, $\overline{\sigma_c}$ can be calculated, of which only the total average cross section can be compared with measurements.
- In OMP one uses transmission coefficients $T_c = 1 - |\overline{U_{cc}}|^2$
- Average single reaction cross section (Hauser-Feshbach):

$$\overline{\sigma_{cc'}} = \overline{\sigma_{cc}^{se}} \delta_{cc'} + \pi \lambda_c^2 g_c \frac{T_c T_{c'}}{\sum T_i} W_{cc'}$$



Measured reaction yield




Cross sections σ_T , σ_γ , σ_n et σ_f



Compound neutron-nucleus reactions



Nuclear level densities




low-lying levels:
Count levels, all J^π


neutron resonances:
Count levels, selected J^π ,
extract D_0

- All level density models reproduce the low-lying levels and D_0 at S_n


Compound nucleus reactions



Compound nucleus reactions




Count the number of levels
in the energy interval → level density




Level densities: the level spacing D_0

- The level spacing D_0 at the neutron binding energy is a crucial input parameter for calibrating level density models.
Level density: $\rho = 1/D$.
- D_0 is the spacing between levels excited by neutrons on nuclei bringing in zero orbital momentum (s-wave resonances).
- Spacings from higher orbital momentum are equally important, but in general much more affected by missing levels.
- Problems concerning the determination of D_0 :
 - spin and parity assignment of levels
 - corrections for missing levels (which are not observed experimentally)


Level spacing D_0



Level spacing D_0



Level spacing D_0



Level density basics

Level density definition:
$$\rho(U, J, \pi) = \frac{\partial N(U, J, \pi)}{\partial E}$$

Simplify, use factorization:
$$\rho(U, J, \pi) = \rho_U(U) \times \rho_J(J) \times \rho_\pi(\pi)$$

• parity distribution:
$$\rho(\pi^+) = \rho(\pi^-) = \frac{1}{2}$$


• spin distribution:
$$\rho(J) = \exp\left(-\frac{J^2}{2\sigma_c^2}\right) - \exp\left(-\frac{(J+1)^2}{2\sigma_c^2}\right)$$

• energy distribution:
(constant temperature)
$$\rho(U) = \frac{1}{T} \exp\left(-\frac{U - U_0}{T}\right)$$


Many more sophisticated models, especially for $\rho(U)$.

Level density by counting levels: staircase plot

$$D(\ell = 0) = D_0 = \Delta E / N \qquad \rho(\ell = 0) = N / \Delta E$$



Level density by counting levels: missing levels



Level density from resonance positions

- Other information needed to estimate the number of missing levels.
- Use the properties of the statistical model of the nucleus to find missing levels.
Works for medium and heavy nuclei.

What is the statistical model for a nucleus?


- Neutron resonances correspond to states in a compound nucleus, which is a nucleus in a highly excited state above the neutron binding energy.
- The compound nucleus corresponds to a very complex particle-hole configuration.
→ **Gaussian Orthogonal Ensemble (GOE)**
- The transition probability between two levels is related to the matrix elements of the interaction between two levels.
- Matrix elements (amplitudes γ) are Gaussian random variables with zero mean. Observables are widths $\Gamma \sim \gamma^2$.

The statistical model

The nucleus at energies around S_n can be described by the **Gaussian Orthogonal Ensemble (GOE)**



The matrix elements governing the nuclear transitions are random variables with a Gaussian distribution with zero mean.

- **Consequences:**
 - The partial widths have a **Porter-Thomas** distribution.
 - The spacing of levels with the same J^π have approximately a **Wigner** distribution.





Chi-square distribution

- If ν random variables x_i have independent Gaussian distributions, $z = \sum x_i^2$ has a chi-square distribution with ν degrees of freedom.



Chi-square distribution

- If ν random variables x_i have independent Gaussian distributions, $z = \sum x_i^2$ has a chi-square distribution with ν degrees of freedom.



- neutron widths $\nu = 1$
- radiation widths $\nu = \text{large number}$
- fission widths $\nu \sim 4$

Chi-square distribution

$$x = \frac{\gamma^2}{\langle \gamma^2 \rangle} \quad P_{\text{PT}}(x) = \frac{1}{\sqrt{2\pi x}} \exp\left(-\frac{x}{2}\right)$$

For neutron widths (s-waves), use the effective reduced neutron width

$$\Gamma_n^0 = \Gamma_n / \sqrt{E}$$

and

$$x = \frac{g\Gamma_n^0}{\langle g\Gamma_n^0 \rangle}$$

Chi-square distribution

$$x = \frac{\gamma^2}{\langle \gamma^2 \rangle} \quad P_{\text{PT}}(x) = \frac{1}{\sqrt{2\pi x}} \exp\left(-\frac{x}{2}\right)$$

For neutron widths (s-waves), use the effective reduced neutron width

$$\Gamma_n^0 = \Gamma_n / \sqrt{E}$$


and

$$x = \frac{g\Gamma_n^0}{\langle g\Gamma_n^0 \rangle}$$

and for easy handling use


$$\int_{x_t}^{\infty} P_{\text{PT}}(x)$$

Neutron widths





Neutron width distribution

$$N(x_t) = N_0 \int_{x_t}^{\infty} P_{PT}(x) dx = N_0(1 - \text{erf}\sqrt{x_t/2})$$





Example ^{61}Ni





Example ^{236}U




The statistical model


The nucleus at energies around S_n can be described by the **Gaussian Orthogonal Ensemble (GOE)**

The matrix elements governing the nuclear transitions are random variables with a Gaussian distribution with zero mean.

- **Consequences:**
 - The partial widths have a **Porter-Thomas** distribution.
 - The spacing of levels with the same J^π have approximately a **Wigner** distribution.



Spacing distribution of two consecutive levels



Evaluated nuclear data libraries

Libraries

- JEFF - Europe
- JENDL - Japon
- ENDF/B - US
- BROND - Russia
- CENDL - China

Common format:

ENDF-6

Contents:

Data for particle-induced reactions (neutrons, protons, gamma, other)
but also radioactive decay data

Data are identified by “materials”
(isotopes, isomeric states, (compounds))

ex. ^{16}O : mat = 825
natV: mat = 2300
 $^{242\text{m}}\text{Am}$: mat = 9547

Files for a material

from report ENDF-102

- 1 General information
- 2 Resonance parameter data
- 3 Reaction cross sections
- 4 Angular distributions for emitted particles
- 5 Energy distributions for emitted particles
- 6 Energy-angle distributions for emitted particles
- 7 Thermal neutron scattering law data
- 8 Radioactivity and fission-product yield data
- 9 Multiplicities for radioactive nuclide production
- 10 Cross sections for photon production
- 12 Multiplicities for photon production
- 13 Cross sections for photon production
- 14 Angular distributions for photon production
- 15 Energy distributions for photon production
- 23 Photo-atomic interaction cross sections
- 27 Atomic form factors or scattering functions for photo-atomic interactions
- 30 Data Covariances obtained from parameter covariances and sensitivities
- 31 Data covariances for nubar
- 32 Data covariances for resonance parameters
- 33 Data covariances for reaction cross sections
- 34 Data covariances for angular distributions
- 35 Data covariances for energy distributions
- 39 Data covariances for radionuclide production yields
- 40 Data covariances for radionuclide production cross sections


Example: part of an evaluated data file

Z and A values	nuclear mass		formalism flag	number of resonances	material number	MF number	MT number
7.919700+4	1.952740+2	0	0	1	07925	2151	1
7.919700+4	1.000000+0	0	0	1	07925	2151	2
1.000000-5	5.000000+3	1	2	0	07925	2151	3
1.500000+0	9.800000-1	0	0	1	07925	2151	4
1.952740+2	0.000000+0	0	0	1578	2637925	2151	5
-3.380000+1	2.000000+0	2.562000-1	1.562000-1	1.000000-1	0.000000+07925	2151	6
4.906000+0	2.000000+0	1.377000-1	1.520000-2	1.225000-1	0.000000+07925	2151	7
4.645000+1	1.000000+0	1.241300-1	1.300000-4	1.240000-1	0.000000+07925	2151	8
5.810000+1	1.000000+0	1.164000-1	4.400000-3	1.120000-1	0.000000+07925	2151	9


Labels for columns (pointing to corresponding values in the table):

- Z and A values
- nuclear mass
- formalism flag
- number of resonances
- material number
- MF number
- MT number
- resonance energy
- spin
- total width
- neutron width
- gamma width
- fission width
- line number


The library JEFF-3.1




Measured reaction yield




Measured reaction yield



Measured reaction yield



Measured reaction yield



Further Reading

Books/articles

- K. S. Krane, *Introductory Nuclear Physics*, Wiley & Sons, (1988).
- G. F. Knoll, *Radiation Detection and Measurement*, Wiley & Sons, (2000).
- P. Reus, *Précis de neutronique*, EDP Sciences, (2003).
- J. E. Lynn, *The Theory of Neutron Resonance Reactions*, Clarendon Press, Oxford, (1968).
- F. Fröhner, *Evaluation and analysis of nuclear resonance data*, JEFF Report 18, OECD/NEA (2000).
- C. Wagemans, *The Nuclear Fission Process*, CRC, (1991).
- A. M. Lane, R. G. Thomas, “R-matrix theory of nuclear reactions”, *Rev. Mod. Phys.* **30** (1958) 257.
- G. Wallerstein, et al., “Synthesis of the elements in stars: forty years of progress”, *Rev. Mod. Phys.* **69** (1997) 995.
- D. Cacuci (ed.), *Handbook of Nuclear Engineering*, Springer (2010).

Internet sites

www.nea.fr
www.nndc.bnl.gov
www.iaea.org
www-instn.cea.fr
www.cern.ch/ntof
www.irmm.jrc.be